

# LSST

## Large Synoptic Survey Telescope

## Active Galaxy Science with the LSST

**W.N. Brandt (PSU), S.F. Anderson (UW), D.R. Ballantyne (UA), A.J. Barth (UCI), R.J. Brunner (UIUC), J. Carson (SLAC), G. Chartas (PSU), W.H. de Vries (LLNL), M. Eracleous (PSU), R.F. Green (LBTO), G.M. Madejski (SLAC), D.P. Schneider (PSU), O. Shemmer (PSU), H.A. Smith (CfA), P.S. Smith (UA), D.E. Vanden Berk (PSU) - The LSST AGN Science Collaboration**


The LSST will provide great advances in our understanding of the demography and physics of active galactic nuclei (AGNs), owing to its unprecedented combination of solid-angle coverage, photometric and astrometric accuracy, sensitivity, broad wavelength coverage, and time sampling. Well-defined, large (20-80 million objects) samples of AGNs at  $0 < z < 7.5$  can be constructed via four approaches: location in color-color space, variability, lack of proper motion, and matching to multiwavelength data. The samples will allow studies of the AGN luminosity function and AGN clustering down to Seyfert luminosities out to  $z \sim 6$ . The time baseline (minutes to years), coupled with the large sample size, will produce a data set that can be used to address the physics of the AGN accretion process, allowing insights into the fueling of AGNs by stellar tidal disruptions and the lifetime of AGNs. Comparison of LSST data with archival X-ray and infrared fields will be the first stage of follow-up for about a million AGNs, many heavily obscured by gas and dust.

### AGN Selection and Yields

The LSST can select AGNs from  $z \sim 0-7.5$  using four effective methods:


- Location in *ugrizy* color-color space. This has been a standard method of AGN selection in wide-field optical surveys, such as the Sloan Digital Sky Survey (SDSS).
- Lack of proper motion. The  $3\sigma$  upper limit on proper motion for the full LSST survey will be a few mas per year.
- Variability on timescales ranging from minutes to years.
- Multiwavelength matching using data from *Chandra*, *XMM-Newton*, *Spitzer*, *GLAST*, and other missions.

In contrast to current wide-field optical surveys, the LSST will find moderate-luminosity, typical AGNs in the high-redshift universe. After 10 years observations, the LSST should deliver AGN sky densities of  $\sim 1000-4000 \text{ deg}^{-2}$ . This is competitive with the deepest pencil-beam multiwavelength AGN surveys to date, such as the Chandra Deep Fields. **The total LSST AGN yield should be  $\sim 20-80$  million AGNs**, which is two orders of magnitude larger than the current AGN census.



### AGNs at the End of the Dark Ages


The LSST can find many AGNs at the highest redshifts using color-selection methods. **About  $\sim 200-1000$  AGNs at  $z \sim 6.5-7.5$  should be found as z-band dropouts** (i.e., detected only in the y-band). These AGNs will have optical luminosities down to  $\sim 10^{44} \text{ erg s}^{-1}$ . By searching for X-ray sources that lack detections in any of the LSST filters, it should be possible to find AGNs at  $z > 7.5$ .



**Fig. 1 - Left:** LSST *z-y* vs. *y* color-magnitude diagram, showing the expected allowable region for redshift  $z \sim 7$  quasar candidates (cyan). The region limits are defined by the  $3\sigma$  *y*-magnitude detection limits, and  $2\sigma$  *z*-magnitude detection limits, as a function of the number of co-added 15 s exposures. An object is considered a redshift  $z \sim 7$  candidate if it is detected at the  $> 3\sigma$  level in the *y*-band, and does not exceed the *z* band  $2\sigma$  detection limit. The *y*-magnitude detection limits reached in a given number of exposures are shown by red dots, and labeled by the number of exposures. The *y*-magnitudes and *z-y* color limits are shown for simulated redshift  $z=7$  quasars at three different 5100 Angstrom continuum luminosities. The open stars show the minimum *z-y* color limits required for a single 15 s exposure, and the ends of the arrows show the limits for 2000 exposures (the full extent of survey). A quasar at redshift 7 with an optical luminosity of  $10^{45} \text{ erg s}^{-1}$  can be detected in about 30 exposures, while quasars with optical luminosities exceeding  $10^{46} \text{ erg s}^{-1}$  can be detected in a single exposure. The expected *z-y* colors of stars (green) and redshift  $< 6$  quasars (blue), based on results from the SDSS, are shown for comparison. The minimum *z-y* colors lie well above the star and low-redshift quasar color distributions. **Right:** Track of a quasar with an optical luminosity of  $10^{46} \text{ erg s}^{-1}$  on the *z-y* vs. *z-z* color-color diagram. Note that such quasars can be easily separated from the loci of stars (black) and lower redshift quasars (colored).

### AGN Variability Studies

We use a parameterized description of AGN variability from the SDSS (extrapolated to fainter apparent magnitudes) to estimate the fraction of AGNs that may be detected as significantly variable by the LSST. We assume that the  $10\sigma$  detection limit in a single-epoch LSST image is  $i \sim 24$ , and calculate the magnitude difference at which only 1% of the non-variable stars will be flagged as variable candidates due to measurement uncertainty. The probability that the single-band rms magnitude difference of an AGN will exceed this value - and will therefore be flagged as a variable candidate - depends upon redshift, luminosity, observed time interval, and the number of observing epochs.





**Fig. 2 -** The probability of detecting an AGN as variable as a function of redshift and absolute magnitude. **Left:** Two epochs separated by 30 days. **Right:** 12 epochs spanning 360 days. Nearly all of the AGNs between the limiting apparent magnitudes would be detected as variable after 1 year.

Even with only two epochs separated by 30 days, a large fraction of AGNs will be detected as variable objects. The fraction of AGNs detected depends strongly on absolute magnitude at each redshift. After 12 epochs with a total time lag of 360 days, **nearly all of the AGNs to a limiting apparent magnitude of  $\sim 24$  will be detected as variable**. The detection fraction will increase as the number of epochs increases, and the use of all six bands will improve the detection fraction even further.

### Stellar Tidal Disruptions

Transient outbursts from galactic nuclei are likely caused by inevitable fueling events of nuclear supermassive black holes when a star, planet, or gas cloud is tidally disrupted and partially accreted. The LSST will greatly extend the limited current studies of such events and will provide a definitive determination of the rate of optical outbursts throughout the Universe. It will explore promising "discovery space" by looking for outbursts with lower luminosities, higher redshifts, and different timescales. Outburst rates will be constrained as a function of galaxy type and nuclear activity. Furthermore, LSST data processing will enable rapid identification of nuclear outbursts, so that **intensive spectroscopic and multiwavelength follow-up studies will be possible while the event is in progress**. Aside from their innate interest, these LSST results will be relevant to the planning of missions such as the Black Hole Finder Probe and LISA.



The effort to build the Large Synoptic Survey Telescope is overseen by the LSST Corporation, a non-profit 501(c)(3) corporation formed in 2003, with headquarters in Tucson, AZ. The LSST research and development effort is funded in part by the National Science Foundation under Scientific Program Order No. 9 (AST-0551161) through Cooperative Agreement AST-0132798. Additional funding comes from private donations, in-kind support at Department of Energy laboratories and other LSST Institutional Members.

Research Corporation • University of Arizona • NOAO • University of Washington • Brookhaven National Laboratory (BNL) • Harvard-Smithsonian Center for Astrophysics (CfA) • Johns Hopkins University • Las Cumbres Observatory, LLC • Lawrence Livermore National Laboratory (LLNL) • Stanford Linear Accelerator Center (SLAC) • Stanford University Kavli Institute for Particle Astrophysics (KIPAC) • University of Pennsylvania • The Pennsylvania State University • University of California, Davis • University of California, Irvine • University of Illinois at Urbana-Champaign • Google Inc.